



Salmon & Trout Conservation

The impact of excess fine sediment on invertebrates and fish in riverine systems

Literature Review

I.	Introduction.....	2
II.	Effects on Invertebrates.....	2
III.	Effects on Fish.....	4
IV.	Ecosystem-Scale Effects.....	7
V.	Current Management.....	8
VI.	Concluding Remarks.....	10
VII.	References.....	12

I. Introduction

At a global scale, suspended solid (SS) concentrations in many rivers have dramatically changed in recent years (Walling, 2006). Existing evidence indicates that natural sediment loadings have been substantially exceeded in many catchments in the UK, particularly since World War II (Evans, 2006). Sediment loadings delivered to watercourses originate from a number of upstream primary and secondary sediment sources, the main anthropogenic activities increasing sediment supply to watercourses include:

- Changes in agricultural practices; for example, increased areas of arable cultivation, leading to greater areas of bare exposed soil susceptible to erosion by winter rainfall (Greig, *et al.*, 2005), and mechanised farm practises which compact the soil and increase runoff and soil erosion (McMellin *et al.*, 2002; Bilotta, *et al.*, 2007). For instance, sediment-fingerprinting research indicated 61% of the sediment load of the River Tweed in Scotland was derived from arable and pasture top soils (Owen *et al.*, 2000).
- Intensification of agricultural practices; for example, increasing stock density (Heaney *et al.*, 2001) and multiple cropping on arable land.
- Increased bank erosion due to loss of natural hydrology.

Erosion processes and sediment delivery form an integral part of aquatic systems, influencing their geomorphology, habitat distribution and water quality. Aquatic communities are adapted, and hence able to cope with, natural 'baseline' sediment inputs. Healthy freshwater ecosystems require sediment inputs to maintain habitat and nutrient fluxes, but excessive loading can have catastrophic effects on river ecosystem function. The main direct physical effects are reduction in habitat availability and modification of habitat biogeochemical conditions through reduction of oxygen and increased concentrations of toxic compounds (Kemp *et al.* 2011; Jones *et al.* 2012). Sediment suspended in the water column can also cause sublethal effects from turbidity and direct physical damage, particularly to fish species (Wilber & Clarke, 2001).

Sediment in the water column can be measured in three main ways 1) turbidity; the optical (light scattering) property of the water, 2) total SS; direct measurement of particulate weight present in a volume of water and 3) water clarity; also an optical property of water. Deposited sediment can also be measured using sediment traps. Despite this, there is a distinct lack of SS monitoring data from around the UK, mainly due to cost implications but also a historic perception that other water quality parameters were of greater importance.

II. Effects on Invertebrates

The negative impacts of high and persistent sediment loads on invertebrate assemblages and abundances are well documented with Ephemeroptera, Plecoptera, Trichoptera (EPT) taxa exhibiting the greatest negative response to increased sediment. Sediment can trigger invertebrate decline in various ways including; scour damage, burial of heavy or immobile species, the clogging of gills or feeding structures, and reduction in interstitial habitat and primary production (Newcombe and MacDonald, 1991) (Fig. 1). Much of the recent literature has explored the effects of excessive sediment deposition on invertebrates from physical and biological perspectives, looking beyond the focus on reduced total abundance and taxonomic richness

found in older papers (Richards & Bacon, 1994; Shaw & Richardson, 2001; Zweig & Rabeni, 2001). Changes in invertebrate life has trophic implications for fish species, particularly juvenile salmonids, so understanding and controlling the impacts of sedimentation is crucial to maintain ecological fitness of river systems.




Figure 1: Summary diagram illustrating the direct and indirect mechanisms by which fine sediments impact upon macro-invertebrates (Jones *et al.* 2012)

Biological changes

A study by Buendia *et al.* (2013) assessed evidence of patterns in assemblage structure and functional traits of benthic invertebrates in response to excessive fine sediment deposition. Invertebrate groupings in high sediment areas were only a subset of the groups found in locations with minimal fine sediment and these invertebrates had biological traits that favoured resistance and resilience to fine sediment, such as shorter life cycles and smaller body sizes. Descloux *et al.* (2014) similarly examined changes in biological characteristics caused by colmation in benthic and hyporheic invertebrate assemblages. In the benthic zone colmation significantly modified eight invertebrate trait profiles and in the hyporheic zone it significantly modified five. As found in the previous paper, trait selection in benthic invertebrates leaned more towards resistance or resilience capacities of species and features related to physiological and trophic functions. As only morphological traits were modified in hyporheic zone invertebrates, it was concluded that colmation is biologically more selective on benthic than hyporheic assemblages as benthic invertebrates have less adaptations to cope with excessive sedimentation.

Behavioural/Habitat changes

Vadher *et al.* (2015) demonstrated via lab experiments that the addition of fine sediment inhibited movement of *Gammarus pulex* into sediment refuges during exposure to water column disturbances. In the control 93% of the individuals moved into subsurface sediments when the water level was reduced. This was reduced to

0% when as little as 20% fine sediment was added. This physical barrier means larger macroinvertebrates cannot protect themselves from stressors such as floods and streambed drying. Reduced availability of refuges caused by fine sediment can also increase the density of invertebrate drift, as invertebrates have nowhere to hide and become forced to leave the river substrate to prevent smothering. Benthic invertebrate drift rates have shown to increase in SS concentration as little as 8mg/l (Rosenberg and Wiens, 1978) and population size has been shown to reduce by 77% when exposed to 62 mg/l of SS for 100 days (Wagener and LaPerriere, 1985). A 40% reduction in stream invertebrate species diversity has been recorded in response to prolonged SS concentrations of 133 mg/l over the period of a year (Nuttall and Bielby, 1973). Molinos & Donohue (2009) examined the response of 4 common macroinvertebrate taxa to different doses and exposures of sedimentation. They found that invertebrate drift was strongly affected by exposure time and all taxa exhibited statistically significant responses within the first day under all exposure concentrations. Larsen & Ormerod (2010) similarly found that even very small, short-term increases in fine sediment reduce benthic density by promoting invertebrate drift. When sediment was added drift density overall increased by 45% with the strongest contributors being mayflies (*Baetis rhodani*, *B. muticus* and *Ecdyonurus* spp.), helodid beetles and simuliid/chironomid dipterans.

III. Effects on Fish

Excessive fine sediment, in suspension or deposited, can have damaging impacts on all life stages of fish, particularly salmonids. This has been worsened for salmonids by a shift in the timings of arable cultivation in the UK, from spring to autumn sown cereals, which now coincides with their egg incubation times (Collins and Walling, 2007; Collins *et al.*, 2008). The effect on ecosystems will, however, depend on several key factors, including: the concentration of fine sediment in suspension; the duration of exposure to fine sediment; and the sediment chemical composition and particle size (Bilotta and Brazier, 2008). This makes determining the impact of SS on fauna and flora difficult to generalise, quantify and compare. However, generic consequences of increased SS concentrations in the water column for fish can include:

Mortality

The relationship between SS concentration and direct mortality is highly complex. The effect of sediment on fish will vary depending on life stage, time of year, size of fish, composition of sediment and the availability of off-channel habitat (Bash *et al.*, 2001), as well as the exposure magnitude, duration and frequency (Servizi and Martens, 1992). For example, in a review of published critical SS concentration thresholds based on dose-response experiments examining impaired growth, reduced feeding and mortality, Berry *et al.* (2003) reported ranges of 27-80,000 mg l⁻¹ for mollusca and 4-330,000 mg l⁻¹ for various fish species. Such ranges in the severity of effect of SS concentration are a function of associated stressors including sediment particle size, species life history stage, temperature, the presence of sediment-associated contaminants and sediment load duration (Swietlik *et al.*, 2003). Due to the complex interaction of such stressors, it is unlikely that a comprehensive list of genus-based critical SS concentration targets can be developed in the short-term (USEPA, 2003).

Reduction in suitable spawning habitat and declines in egg/early life stage success

Effects of excessive deposition of fine sediment on salmonid spawning success and egg survival have been well documented over the years. It has been proved in a vast plethora of literature that infiltration of fine sediment limits success of eggs hatching through the reduction of gravel permeability and oxygen availability (Ingendahl, 2001; Greig *et al.* 2007; Schindler Wildhaber *et al.* 2014). Salmonid eggs (as well as many cyprinid fish and lamprey eggs) require a well-oxygenated environment during the embryonic development stage, so eggs are laid in permeable gravel beds with interstitial pore spaces, which allow the passage of oxygenated water. Excess fine sediment in the water, when deposited, can clog these interstitial pores, obstructing the circulation of oxygenated water, which reduces egg survival (Carling, 1984; Magee *et al.* 1996). Salmonid egg mortality of between 98-100% has been recorded within spawning gravels experiencing high siltation loads (Turnpenny and Williams, 1980). The effect is particularly damaging when sediments contain a high organic component, as its subsequent decomposition also reduces oxygen from the water (Rubin, 1998). Excess deposited sediment can also reduce interstitial and hyporheic (region beneath streambed) flow velocities (Chapman, 1988; Acornley and Sear, 1999). This decreases the natural flushing process, which removes the harmful metabolic waste excreted by the embryos (Burkhalter and Kaya, 1975). SS can also be damaging to fish species, such as perch and roach, depositing eggs on macrophytes, as silt particles can adhere to the eggs and reduce oxygen and carbon dioxide exchange (Stuart, 1953).




Figure 2: The three specific mechanisms by which fine sediment accumulation restricts O₂ availability to incubating salmonid embryos (Greig *et al.* 2005)

Fine sediment can also exert sub-lethal effects on fish fry including: delaying emergence by trapping fry in interstitial pores (Phillips *et al.*, 1975); and premature hatching of smaller and poorer quality fry, due to exposure to low dissolved oxygen concentrations (Alderdice *et al.*, 1958; Mason, 1969). Researchers have found a relationship between fine sediment (less than 0.850 mm) and fry survival, with decreasing survival of up to 3.4% for each 1% increase in fine sediment (Cederholm *et al.*, 1981). Louhi *et al.* (2011) showed that brown trout (*Salmo trutta*) fry experienced predator-masking effects associated with high sedimentation. Control fish tended to postpone fry emergence when exposed to predator odour, whereas fish in the high-sedimentation treatment showed no response to predators. This indicates that in high sediment environments predation is a bigger risk to fry survival. The study also confirmed that in high sediment environments trout fry were emerging earlier with larger yolk sacs. As these fry emerged prematurely to

escape the suboptimal conditions associated with increased sedimentation it is highly probable that they will be poor swimmers and thus more vulnerable to predation and downstream displacement than fully developed fry. Bowerman *et al.* (2014) found the same phenomenon occurred in bull trout (*Salvelinus confluentus*) fry; they too were emerging from the incubation environment at much earlier development stages when sediment percentages were higher. This work also highlighted later life implications for premature fry that do survive to adulthood, as early emergence can lead to fitness reduction in later life stages. The deposition of sediment on the riverbed also changes and degrades physical habitat for bottom dwelling fish and fry, leading to lower fry and parr density (Lisle and Lewis, 1992). The sediment fills in the spaces between substrate particles, creating a smoother riverbed (Diplas and Parker, 1992). This, not only reduces the available habitat complexity and availability, but also increases water velocity and the need for shelter from the water current (Richardson and Jowett, 2002). Sedimentation also decreases habitat connectivity (Cohen, 1995) and heterogeneity (Boles, 1981).

Gill irritation/trauma

Fish gills are delicate and easily damaged by abrasive sediment particles. Fish species have been found with increasing levels of deformities, eroded fins, lesions, tumours, gill flaring and 'coughing', all related to increasing SS in the water column (Berg, 1982; Schleiger, 2000).

Altered blood physiology

Research has show increases in plasma glucose (Servizi and Martens, 1987), blood sugar levels (Servizi and Martens, 1992) and plasma cortisol (Redding *et al.*, 1987) in fish species exposed to fine sediment. These are all indicators of stress, and can impact physiological systems, reduce growth and decrease immunological competence against other stressors, such as disease. In a study by Sutherland & Meyer (2007) all three life stages of the spotfin chub (*Erimonax monachus*) and whitetail shiner (*Cyprinella galactura*) showed increases in immunoreactive corticosteroid (stress) levels when exposed to moderate sediment concentrations. This study showed that fish species experience physiological stress as a result of suspended sediment regardless of their life stage. Respiratory impairment caused by sediment in the gills was suggested to be the main factor instigating the stress response in the spotfin chubs. Stress to salmonids can also affect the smoltication process, leading to decreased osmoregulatory ability, impaired migrations and a reduction in early marine survival (Wedemeyer and McLeay, 1981).

Altered movement/swimming performance

Migrating fish species, such as salmonids, are commonly known to migrate through high SS concentrations in estuaries. However, like other containments, exposure time is a key element in the impact of SS as well as concentration (Newcombe and MacDonald, 1991). This means high exposure rates to sediment loads can halt fish migration, particularly upstream. Fish are known to exhibit avoidance reactions and move away from the vicinity of adverse sediment conditions, if refuge conditions are present (Sigler *et al.*, 1984; Bash *et al.*, 2001). The effects of suspended sediment on swimming ability on juvenile brown trout (*Salmo trutta*) and rainbow trout (*Oncorhynchus mykiss*) were explored by Berli *et al.* (2014). Both species experienced a decrease in swimming performance as turbidity increased, but rainbow trout were impaired to a greater extent. This was attributed to impairment in the ability of the fish to utilise anaerobic metabolic pathways in high sediment environments. The authors concluded that the ability of salmonids to maintain swimming

performance is hindered when fish are exposed to environmentally relevant, suspended sediment-generated turbidities.

Changed foraging behaviour and reduced territoriality

Turbidity can reduce feeding rates, and affect prey selection and prey abundance. This is particularly significant for visual feeders, such as salmonids, where SS can reduce the effectiveness of them obtaining food (Berg, 1982). However, research also suggests the turbid-clear water interface may sometimes assist feeding, by offering concealment and protection within the turbid water (Scullion and Edwards, 1980). Robertson *et al.* (2007) showed that during autumn initial introduction of sediment increased foraging activity in juvenile Atlantic salmon (*Salmo salar*). It was suggested that the small increase in turbidity provided visual isolation from predators. As sediment levels were raised to over 180 mg/L this foraging activity declined along with a rapid reduction in territorial behaviour. Such responses to increased water turbidity have been shown to instigate increased emigration from preferred habitats as excess sediment makes habitats unsuitable. Although salmon were less responsive to increases in suspended sediment levels in colder water temperatures (winter months) this work highlighted the importance of not assuming salmon are more sediment tolerant at these times. Pulses of turbid water have also been shown to breakdown normal social organisation and territoriality, which can decrease feeding rates and may affect overall growth rates (Berg, 1982).

IV. Ecosystem-scale effects

Community homogenisation

Sedimentation can affect aquatic biota at both a population and community level, and result in homogenisation. This means regionally distinct faunas may be replaced with few invasive and disturbance tolerant species (Walters *et al.*, 2003). This could be a serious threat to biodiversity, both now and in the future, by reducing species' resilience to climate change. Sedimentation can also increase the susceptibility of invasive species such as Canadian pondweed and the common carp, which potentially have major disruptive effects on aquatic ecosystems.

Transfer of pollutants

Fine sediment exerts an important control on the transfer and fate of a wide range of agricultural and industrial contaminants (Warren *et al.*, 2003; Collins *et al.*, 2005). Sediment therefore represents an important vector for contaminants such as phosphorus (Haygarth *et al.*, 2005); heavy metals (Neal *et al.*, 1999) and organic pollutants like sheep dip substances (Long *et al.*, 1998). These associated pollutants can alter species assemblages by poisoning the water system, and accelerating eutrophication. The capacity of fine sediment to bind contaminants can also lead to an increase in the resident times of the pollutants in aquatic systems (Foster and Charlesworth, 1996), thereby increasing exposure times and the opportunity for pollutant remobilisation.

Reduced primary productivity

Suspended solids reduce water clarity and increase turbidity, exerting a negative effect upon primary production. Research suggests in subarctic Alaskan streams concentrations of SS as little as 8mg/l can reduce primary production by 3-13% (Lloyd, 1987), and above 2100 mg/l no primary production can occur (Van Nieuwenhuysse and LaPerriere, 1986).

Depressed oxygen levels in the water

Suspended solids can contribute towards raising the Biological Oxygen Demand (BOD; Petts *et al.*, 2002), and hence lowered oxygen levels potentially to stressful or lethal levels for vulnerable species and life stages.




Figure 3: Effects of high sediment loads on aquatic ecosystems. Rectangles = Physicochemical effects. Ovals = Direct/long-term biological and ecological responses. (Kemp *et al.* 2011)

V. Current Management

Until it was revoked, the Freshwater Fish Directive (FFD) stipulated that suspended solid concentrations should not exceed an annual mean of 25mg/l. However, this standard was not imperative and the recommended figure was simply a guideline. In 2009 the FFD and Shellfish Waters Directive were both revoked, with their key requirements being transferred to the Water Framework Directive (WFD). In 2010,

UKTAG published a note detailing recommendations that should be considered in the transfer, including an environmentally protective solids standard. Application of a standard was deemed necessary to protect shellfish against potential smothering, release of sediment associated pollutants and dissolved oxygen sags related particularly to dredging activity. It was also recommended that suspended solids continue to be monitored due to their ecological significance (UKTAG, 2010). Since the repeal of both Directives, the initial FFD guideline target was lost and no standards for suspended solids under the Water Framework Directive currently exist (Cascade Consulting, Thames Water Utilities, 2011).

In the scientific community attempts have been made to identify target values for both deposited fine sediment and sediment loading, yet the relationship between deposited fine sediment and agricultural sediment pressure is still poorly understood. Collins *et al.* (2007) used a structured modelling methodology to predict the impact of projected structural evolution in agriculture (land use change) and the uptake of sediment mitigation methods due to programmes like the England Catchment Sensitive Farming Delivery Initiative (ECSFDI) on annual mean suspended sediment concentrations across England and Wales by 2015. This work suggested that structural and mitigation work could potentially reduce the national sediment loss from the agricultural sector by 9% by 2015.

Collins and Anthony (2008a) also modelled catchment compliance across England and Wales using the previous FFD guideline standard. The study provided the first national scale assessment of sediment sources for England and Wales under current environmental conditions (year 2000), suggesting that source contributions are in the order: agricultural sector (1929 kt = 76%) eroding channel banks (394 kt = 15%), diffuse urban sources (147 kt = 6%) and point source discharges (76 kt = 3%). A structured regression model was used to convert the predicted total sediment loadings into time-averaged suspended sediment concentrations at national scale. The findings suggested that approximately 83% of the total catchment area of England and Wales appears to require no further reductions in sediment loss to rivers from diffuse agricultural sources for the purpose of meeting 'Good Ecological Status' (GES) as defined by the Water Framework Directive (WFD). It is important to note, however, that the use of the FFD sediment threshold failed to identify catchments across England and Wales where the detrimental impacts of sediment are widely reported e.g. the chalk catchments of southern and eastern England. Chalk catchments are particularly vulnerable to sedimentation due to the lack of any significant flushing effect owing to their baseflow-dominated hydrology.

Naden *et al.* (2016) took these concepts further and analysed instantaneous measurements of deposited fine sediment in 230 agricultural streams across England and Wales in relation to 20 potential explanatory catchment and channel variables. Two main practical recommendations were made from this work regarding fine sediment load targets: 1) the ability of streams to transport/retain fine sediment needs to be taken into account, 2) where agricultural mitigation measures are implemented to reduce delivery of sediment, river management to mobilise/remove fines may also be needed in order to effect an improvement in ecological status in cases where streams are already saturated with fines and unlikely to self-cleanse.

There are, however, serious concerns regarding the use of a single global threshold concentration for suspended sediment. This is because of the large variability of effects caused by sediment, the diversity of

habitat and conditions within catchments, the existence of sub-threshold effects on both fish and their supporting ecosystems, and the failure of an annual mean to capture the highly episodic nature of sediment pressures which are focused during flood events. To address this alternative sediment targets were proposed for England and Wales using an alternative sediment target scheme (Cooper *et al.*, 2008). This was based on nationally extrapolated suspended sediment yields and uses the lower quartile of the measured ranges for catchment types to represent potential targets and the upper quartiles as critical thresholds. These tentative targets were intended for use in the identification of thresholds from a local perspective. Collins and Anthony (2008b) used a structured modelling framework taking explicit account of sediment sources derived from different societal sectors to assess catchment compliance at national scale across England and Wales using these alternative sediment targets. This work successfully identified catchments where negative sediment impacts on fish are being reported.

The use of sediment yields to represent sediment targets is undermined by a number of problems. Since suspended sediment fluxes represent the aggregate of sediment delivery, their utility is best found in helping to define overall catchment response to environmental pressures as opposed to ecological impacts. Reliable coupling of sediment loadings to ecological impacts requires understanding of additional metrics such as sediment deposition and flushing and sediment grain size characteristics. It also important to highlight that all modelling data is based by common consensus on inadequate knowledge of all pathways and adequate monitoring data assumptions and therefore should not be used in isolation, but as part of an integrated modelling and monitoring approach, to help manage uncertainty and ground truth results. Anecdotal evidence from stakeholders on the impacts of fine sediments upon ecosystem can also provide important insight, and therefore should not be ignored.

VI. Concluding Remarks

Understanding and monitoring sedimentation pressure is key to ensuring the delicate ecological mechanisms of riverine ecosystems are preserved. Natural sediment pressures within river systems vary dramatically depending on catchment topography, geology, vegetation, local climate and land use (Hicks and Griffiths, 1992). It is now accepted that excess sediment can cause deterioration in water quality and aquatic biodiversity (Collins *et al.*, 2008). The evidence here highlights the threats our aquatic fauna and flora face because of excess fine sediment pressures. The WFD objective of GES cannot be achieved without addressing this important pressure. Given the problems associated with using the FFD threshold or the alternative sediment yield based target scheme, urgent action is required to identify more meaningful revised sediment targets for England and Wales. Revised targets must take more explicit account of the impacts of sediment on aquatic ecology and should be developed in a catchment-specific manner (Collins and McGonigle, 2008). A generic measurement of sedimentation is not reliable; therefore management should focus on the river basin scale to ensure source control, taking more account of observed impacts rather than modelled inputs. Preventing further damage to river habitats and associated species requires catchment-scale, holistic management, involving the cooperation and regulation of all land users. Managing excess sediment requires prevention and restoration measures, all of which require sound understanding of the key sources (Collins and Walling, 2004) and appropriate monitoring to gauge catchment compliance against revised and improved catchment-specific sediment targets. In order for sediment management to progress in

England and Wales, better-informed sediment targets, and replicable monitoring methods are urgently required for compliance testing.

For further information please contact Lauren Mattingley (Scientific Assistant for S&TC) at:

lauren@salmon-trout.org

© 2017 SALMON & TROUT CONSERVATION, ALL RIGHTS RESERVED

VII. References

- Acornley, R.M. and Sear, D.A. (1999). Sediment transport and sedimentation of Brown trout (*Salmo trutta* L) spawning gravels in chalk streams. *Hydrol. Process.* **13**: 447-458
- Alderdice, D.W., Wickett, W.P. and Brett, J.R. (1958). Some effects of exposure to low dissolved oxygen levels on Pacific salmon eggs. *J. Fish. Res. Board Can.* **15**: 229-250
- Bash, J., Berman, C. and Bolton, S. (2001). Effects of Turbidity and Suspended Solids on Salmonids. Centre for Streamside Studies, University of Washington
- Berg, L. (1982). The effect of exposure to short-term pulses of suspended sediment on the behaviour of juvenile salmonids. In: Hartman, G.F. (Ed). Proceedings of the Carnation Creek workshop: a ten-year review. Department of Fisheries and Oceans, Pacific Biological Station, Nanaimo, Canada
- Berli, B.I. et al., 2014. Acute exposure to a common suspended sediment affects the swimming performance and physiology of juvenile salmonids. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **176**: 1–10
- Berry, W., Rubinstein, N., Melzian, B. and Hill, B. (2003). The biological effects of suspended and bedded sediment (SABS) in aquatic systems: a review. US Environment Protection Agency, National Health and Environmental Health Effects Laboratory, Rhode Island, Internal Report, 58pp.
- Bilotta, G.S. and Brazier, R.E. (2008). Understanding the influence of suspended solids on water quality and aquatic biota. *Water Res.* doi: 10/1016/j.watre.2008.03.018.
- Bilotta, G.S., Brazier, R.E. and Haygarth, P.M. (2007). The impacts of grazing animals on the quality of soils, vegetation, and surface waters in intensively managed grasslands. *Advances in Agronomy* **94**: 237-280
- Boles, G.R. (1981). Macro-invertebrate colonisation of replacement substrate below a hypolimnial release reservoir. *Hydrobiologia* **78**: 133-146.
- Bowerman, T., Neilson, B. & Budy, P., 2014. Effects of fine sediment, hyporheic flow, and spawning site characteristics on survival and development of bull trout embryos. *Canadian Journal of Fisheries and Aquatic Sciences*, **71**(7): 1059–1071
- Buendia, C. et al. (2013). Detecting the structural and functional impacts of fine sediment on stream invertebrates. *Ecological Indicators*, **25**: 184–196
- Burkhalter, D.E. and Kaya, C.M. (1975). Effects of prolonged exposure to ammonia on fertilised eggs and sac fry of Rainbow trout (*Salmo gairdneri*). *Trans. Am. Fish. Soc.* **106**: 470-475
- Carling, P.A. (1984). Deposition of fine and coarse sands in an open-work gravel bed. *Can. J. Fish. Aquat. Sci.* **41**: 263-270
- Cascade Consulting, Thames Water Utilities (2011) *Technical Appendix C, Water Quality Assessment*
- Cederholm, C.J., Reid, L.M. and Salo, E.O. (1981). The effects of logging road landslide siltation on the salmon and trout spawning gravels of Stequaleho Creek and the Clearwater River Basin, Jefferson County, Washington, 1972-1978. FRI-UW-7915. Fisheries Research Institute, University of Washington, Seattle
- Chapman, D.W. (1988). Critical review of variables used to define effects of fine redds of large salmonids. *Trans. Am. Fish. Soc.* **117**: 1-21
- Cohen, A.S. (1995). Paleoecological approaches to the conservation biology of benthos in ancient lakes: a case study from Lake Tanganyika. *Journal of the North American Benthological Society* **14**: 654-668
- Collins, A.L. and Anthony, S.G. (2008a). Assessing the likelihood of catchments across England and Wales meeting 'good ecological status' due to sediment contributions from agricultural sources. *Environmental Science and Policy* **11**: 163-170.
- Collins, A.L. and Anthony, S.G. (2008b) Predicting sediment inputs to aquatic ecosystems across England and Wales due to current environmental conditions. In press: *Applied Geography*
- Collins, A.L., Anthony, S.G., Turner, T. and Hawley, J. (2007). Predicting the impact of projected change in agriculture by 2015 on annual mean fluvial suspended sediment concentrations across England and Wales. In: *Water Quality and sediment behaviour of the future: predictions for the 21st century* (pp 28-37), International Association of Hydrological Sciences Publication No. 314, Wallingford, UK
- Collins, A.L. and McGonigle, D.F. (2008). Monitoring and modelling diffuse pollution from agriculture for policy support: UK and European experience. *Environmental Science and Policy* **11**: 97-101
- Collins, A.L., Walling, D.E. and Peet, L. (2008). Identifying gravel siltation sources. *Salmo Trutta* **11**: 26-28
- Collins, A.L. and Walling, D.E. (2007). Sources of fine sediment recovered from the channel bed of lowland groundwater-fed catchments in the UK. *Geomorphology* **88**: 120-138
- Collins, A.L., Walling, D.E. and Leeks, G.J.L. (2005). Storage of fine-grained sediment and associated contaminants within the channels of lowland permeable catchments in the UK. In: *Sediment Budgets 1* (pp 259-268), International Association of Hydrological Sciences Publication No. 291, Wallingford, UK
- Collins, A.L. and Walling, D.E. (2004). Documenting catchment suspended sediment sources: problems, approaches and prospects. *Progress in Physical Geography* **28**:159-196
- Cooper, D.M., Naden, P., Old, G. & Laize, C. (2008). *Development of guideline sediment targets to support management of sediment inputs into aquatic systems*. Natural England Research Report NERR008. Natural England, Sheffield
- Descloux, S., Detry, T. & Usseglio-Polatera, P (2014). Trait-based structure of invertebrates along a gradient of sediment colmation: Benthos versus hyporheos responses. *Science of the Total Environment*, **466-467**: 265–276
- Diplas, P. and Parker, G. (1992). *Deposition and removal of fines in gravel-bed streams*. In: Billi, P. Hey, R.D., Thorne, C.R. and Tacconi, R. (Eds). Dynamics of gravel-bed rivers. London. John Wiley and Sons
- Evans, R. (2006). Land use, sediment delivery and sediment yield in England and Wales. In: Owens, P.N. and Collins, A.J. (eds), Soil erosion and sediment redistribution in river catchments. CAB International, Wallingford, pp 70-84
- Foster, I.D.L. and Charlesworth, S.M. (1996). Heavy metals in the hydrological cycle: trends and explanation. *Hydrol. Process* **10**: 227-261
- Greig, S., Sear, D. & Carling, P. (2007). A field-based assessment of oxygen supply to incubating Atlantic salmon (*Salmo salar*) embryos. *Hydrological Processes*, **21**(22): 3087–3100
- Greig, S.M., Sear, D.A. and Carling, P.A. (2005). The impact of fine sediment accumulation on the survival of incubating salmon progeny: Implications for sediment management. *Science of the Total Environment* **344**: 241-258
- Haygarth, P.M., Condron, L.M., Heathwaite, A.L., Turner, B.L. and Harris, G.P. (2005). The phosphorous transfer continuum: linking source to impact with an interdisciplinary and multi-scale approach. *Science of the Total Environment* **344**: 5-14
- Heaney, S.I., Foy, R.H., Kennedy, G.J.A., Crozier, W.W. and O'Connor, W.C.K. (2001). Impacts of agricultural on aquatic systems: lesson learnt and new unknowns in Northern Ireland. *Mar. Freshw. Res.* **52**: 151-163
- Hicks, D.M. and Griffiths, G.A. (1992). *Sediment load*. In: Mosley, M. P. (Ed). Waters of New Zealand. Wellington, New Zealand, New Zealand Hydrological Society
- Ingendahl, D. (2001). Dissolved oxygen concentration and emergence of sea trout fry from natural redds in tributaries of the River Rhine. *Journal of Fish Biology*, **58**: 325–341
- Jones, J.I. et al. (2012). The Impact Of Fine Sediment On Macro-Invertebrates. *River Research and Applications*, **28**(8): 1055–1071

- Kemp, P. *et al.* (2011). The impacts of fine sediment on riverine fish. *Hydrological Processes*, **25**(11): 1800–1821
- Larsen, S. & Ormerod, S.J. (2010). Low-level effects of inert sediments on temperate stream invertebrates. *Freshwater Biology*, **55**(2): 476–486
- Lisle, T.E. and Lewis, J. (1992). Effects of sediment transport on survival of salmonids embryos in a natural stream: a simulation approach. *Can. J. Fish. Aquat. Sci.* **49**: 2337–2344
- Lloyd, D.S. (1987). Turbidity as a water quality standard for salmonid habitats in Alaska. *North. Amer. J. Fish. Management*. **7**: 34–45
- Long, J.L.A., House, W.A., Parker, A. and Rae, J.E. (1998). Micro-organic compounds associated with sediments in the Humber rivers. *Science of the Total Environment* **210**: 229–253
- Louhi, P. *et al.* (2011). Does fine sediment constrain salmonid alevin development and survival? *Canadian Journal of Fisheries and Aquatic Sciences*, **68**(10): 1819–1826
- Magee, J.P., McMahon, T.E. and Thurow, R.F. (1996). Partial variation in spawning habitat of cutthroat trout in a sediment-rich stream basin. *Trans. Am. Fish. Soc.* **125**: 768–779
- Mason, J.C. (1969). Hypoxial stress prior to emergence and competition among coho salmon. *J. Fish. Res. Board Can.* **26**: 63–91
- McMellin, G., Walling, D.E. and Nicholls, D. (2002). *Land use and Fisheries*. Report W2-046/TR1. Bristol (UK): Environment Agency
- Molinos, J.G. & Donohue, I. (2009). Differential contribution of concentration and exposure time to sediment dose effects on stream biota. *Journal of the North American Benthological Society*, **28**(1): 110–121
- Naden, P.S. *et al.* (2016). Understanding the controls on deposited fine sediment in the streams of agricultural catchments. *Science of The Total Environment*, **547**: 366–381
- Neal, C., Jarvie, H.P. and Oguchi, T. (1999). Acid-available particulate trace metals associated with suspended sediment in the Humber rivers: a regional assessment. *Hydrological Processes* **13**: 1011–1034
- Newcombe, C.P. and MacDonald, D.D. (1991). Effects of suspended sediments of aquatic ecosystems. *North American Journal of Fisheries Management* **11**: 72–82
- Nuttall, P.M. and Bielby, G.H. (1973). The effect of china-clay wastes on stream invertebrates. *Environ. Pollut.* **5**: 77–86
- Owens, P.N., Walling, D.E., and Leeks, G.L.J. (2000). *Tracing fluvial suspended sediment sources in the catchment of the River Tweed, Scotland, using composite fingerprints and a numerical mixing model*. In: Foster, I.D.L. (Ed). *Tracers in Geomorphology*. Wiley, Chichester: 291–308
- Petts, G.E., Heathcote, J. and Martin, D. (2002). *Urban Rivers: Our Inheritance and Future*. IWA Publishing/ Environment Agency. 117pp
- Phillips, R.W., Lantz, R.L., Claire, E.W. and Moring, J.R. (1975). Some effects of gravel mixtures on emergence of coho salmon and steelhead trout fry. *Trans. Fish. Soc.* **103** (3): 461–466
- Redding, J.M., Schreck, C.B. and Everest, F.H. (1987). Physiological effects on coho salmon and steelhead of exposure to suspended solids. *Trans. Amer. Fish. Soc.* **116**: 737–744
- Richards, C. & Bacon, K.L. (1994). Influence of fine sediment on macroinvertebrate colonization of surface and hyporheic stream substrates. *The Great Basin Naturalist*, **54**(2): 106–113
- Richardson, J. and Jowett, I.G. (2002). Effects of sediment on fish communities in East Cape streams, North Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* **36**: 431–442
- Robertson, M.J., Scruton, D.A. & Clarke, K.D. (2007). Seasonal effects of suspended sediment on the behavior of juvenile Atlantic salmon. *Transactions of the American Fisheries Society*, **136**(3): 822–828
- Rosenberg, D.M. and Weins, A.P. (1978). Effects of sediment addition on macrobenthic invertebrates in a northern Canadian river. *Water Research* **12**: 753–763
- Rubin, J.F. (1998). Survival and emergence of sea trout fry in substrata of different compositions. *J. Fish. Biol.* **53**: 84–92
- Schindler Wildhaber, Y. *et al.* (2014). Artificial steps mitigate the effect of fine sediment on the survival of brown trout embryos in a heavily modified river. *Freshwater Biology*, **59**: 544–556
- Schleiger, S.L. (2000). Use of an index of biotic integrity to detect effects of land use on stream fish communities in wet-central Georgia. *Trans. Am. Fish. Soc.* **129**: 1118–1183
- Scullion, J. and Edwards, R.W. (1980). The effects of coal industry pollutants on the macro-invertebrate fauna of a small river in the South Wales Coalfield. *Freshwater Biol.* **10**: 141–162
- Servizi, J.A. and Martens, D.W. (1992). Sub lethal responses of coho salmon (*Oncorhynchus kisutch*) to suspended sediments. *Can. J. Fish. Aquat. Sci.* **49**: 1389–1395
- Servizi, J.A. and Martens, D.W. (1987). Some effects of suspended Fraser River sediments on sockeye salmon (*Oncorhynchus nerka*). In: Smith, H.D., Margolis, L. and Wood, L.L. (eds). *Sockeye salmon (*Oncorhynchus nerka*) population biology and future management*. *Can. Spec. Publ. Fish. Aquat. Sci.* **96**:254–264
- Shaw, E.A. & Richardson, J.S. (2001). Direct and indirect effects of sediment pulse duration on stream invertebrate assemblages and rainbow trout (*Oncorhynchus mykiss*) growth and survival. *Canadian Journal of Fisheries and Aquatic Sciences*, **58**(11): 2213–2221
- Sigler, J.W., Bjornn, T.C. and Everest, F.H. (1984). Effects of chronic turbidity on density and growth of steelheads and coho salmon. *Transactions of the American Fisheries Society* **113**: 142–150
- Stuart, T.A. (1953). Spawning migration, reproduction and young stages of loch trout (*Salmo trutta*). *Freshw. Salm. Fish. Res.* **5**: 1–39.
- Sutherland, A.B. & Meyer, J.L. (2007). Effects of increased suspended sediment on growth rate and gill condition of two southern Appalachian minnows. *Environmental Biology of Fishes*, **80**(4): 389–403
- Swietlik, W., Berry, W., Gardner, T., Hill, B., Jha, M., Kaufmann, P., Melzian, B., Norton, D., Paul, J., Rubinstein, N., Shippen, R., Spehar, R. (2003). Developing water quality criteria for suspended and bedded sediments (SABS). Potential approaches. US EPA Office of Water, Office of Science and Technology, Health and Ecological Criteria Division, USA
- Turnpenny, A.W.H. and Williams, R. (1982). A conductometric technique for measuring the water velocity in salmonid spawning beds. *Water Res.* **16**: 1383–1390
- UKTAG, 2010. *Water Framework Directive: An approach to the Revoked Directives: – the Freshwater Fish Directive, the Shellfish Directive and the Dangerous substances Directive*
- USEPA. (2003). Developing water quality criteria for suspended and bedded sediments (SABS): potential approaches. US Environment Protection Agency, Office of Water, Office of Science and Technology, Washington DC, USA
- Vadher, A.N., Stubbington, R. & Wood, P.J. (2015). Fine sediment reduces vertical migrations of *Gammarus pulex* (Crustacea: Amphipoda) in response to surface water loss. *Hydrobiologia*, **753**(1): 61–71
- Van Nieuwenhuysse, E.E. and LaPerriere, J.D. (1986). Effects of placer gold mining on primary production in sub-arctic streams of Alaska. *Water Resources Bulletin* **22**: 91–99
- Wagener, S.M. and LaPerriere, J.D. (1985). Effects of placer mining on the invertebrate communities of interior Alaska. *Freshwater Invertebrate Biol.* **4**: 208–214
- Walling, D.E. (2006). Human impact on land-ocean sediment transfer by the world's river. *Geomorphology* **79**: 192–216
- Walters, D.M., Leigh, D.S. and Bearden, A.B. (2003). Urbanisation, sedimentation and the homogenization of fish assemblages in

- the Etyowah River Basin, USA. *Hydrobiologia* **494**: 5-10
- Warren, N., Allan, I.J., Cater, J.E., House, W.A., Parker, A. (2003). Pesticides and other micro-organic contaminants in freshwater sedimentary environments – a review. *Applied Geochemistry* **18**: 159-194
- Wedemeyer, G.A. and McLeay, D.J. (1981). *Methods for Determining the Tolerance of Fishes to Environmental Stressors*. In: Pickering, A.D. (ed). Stress and fish. Academic Press, Toronto, Ont.
- Wilber, D.H. & Clarke, D.G. (2001). Biological Effects of Suspended Sediments: A Review of Suspended Sediment Impacts on Fish and Shellfish with Relation to Dredging Activities in Estuaries. *North American Journal of Fisheries Management*, **21**: 855–875
- Zweig, L.D. & Rabeni, C.F. (2001). Biomonitoring for deposited sediment using benthic invertebrates : a test on 4 Missouri streams. *Journal of the North American Benthological Society*, **20**(4): 643–657